



# The measurement of open apices of teeth to test chronological age of over 14-year olds in living subjects

Roberto Cameriere<sup>a,\*</sup>, Hervoje Brkic<sup>b</sup>, Branko Ermenc<sup>c</sup>, Luigi Ferrante<sup>d</sup>,  
Maja Ovsenik<sup>c</sup>, Mariano Cingolani<sup>a</sup>

<sup>a</sup> *Institute of Forensic Medicine, University of Macerata, Via Don Minzoni 9, 62100 Macerata, Italy*

<sup>b</sup> *School of Dental Medicine, Department of Dental Anthropology, University of Zagreb, Gunduličeva 5, HR-10000 Zagreb, Croatia*

<sup>c</sup> *Institute of Forensic Medicine, Faculty of Medicine, University of Ljubljana, Korytkova 2, 1000 Ljubljana, Slovenia*

<sup>d</sup> *Institute of Biochemical Biotechnologies, Polytechnic University of Marche, Ancona, via Ranieri 65, 60100 Ancona, Italy*

Received 26 November 2006; received in revised form 16 March 2007; accepted 25 April 2007

## Abstract

Age determination in living subjects is a problem of increasing interest in our community, due to the increasing numbers of individuals without identification papers, who have immigrated illegally or committed crimes, and for whom it is necessary to verify whether they have reached the age of 14 years in order to be charged legally.

Although the most widespread methods for age estimation refer to skeletal or dental analysis, these methods do present some drawbacks for identification of the age of 14. The aim of the present study is to discriminate between children who are or are not 14 years of age or older by measuring the open apices of teeth.

We evaluated the OPGs of 447 persons aged between 12 and 16 years, of Italian, Croatian and Slovenian nationality. For each individual, dental maturity was estimated using the number of the seven left permanent mandibular teeth with root development complete, and normalized measurement of the open apices of the third molar.

The results revealed that an individual is considered to be 14 years of age or older if all seven left permanent mandibular teeth have closed apices and the normalized measurement of open apices of the third molar is lower than 1.1.

© 2007 Published by Elsevier Ireland Ltd.

**Keywords:** Forensic odontology; Age estimation; Third molar; Open apices of teeth

## 1. Introduction

The need to determine the age of living individuals is a problem of increasing interest in our community, due to the progressively higher number of persons not in possession of any document of identity or whose birth certificate may be suspected to be wrong, who have immigrated illegally or committed crimes, whose real age must be known in order to decide whether they can be charged, and whether they should be subjected to trial as of age or at least 14 years old. Also in cases of adoption, it is sometimes important to assess age when no birth certificate is available. In the last few years, therefore, forensic medicine has shown increasing interest in this problem

[1–5] and in the reliability of methods for assessing biological age.

During the growth of a person, skeletal, odontological, anthropological and psychological methods allow an approximate assessment of age. Among the methods most frequently used for skeletal assessment are those concerning the left hand–wrist area [6] and FELS [7], which can produce estimates up to the age of 16 years, at which time wrist maturation is complete in 90% of subjects.

Numerous odontological studies have also been carried out to establish age, assessing mineralization within acceptable error limits.

The most common method for dental age assessment was first published by Demirjian et al. [8] and since then odontology has carried out numerous studies in this issue [9].

Nevertheless, to the best of our knowledge, few papers were addressed to assess if an individual is at least 14 years old. Since

\* Corresponding author. Tel.: +39 0733 2582625; fax: +39 0733 2582635.

E-mail address: [r.cameriere@unimc.it](mailto:r.cameriere@unimc.it) (R. Cameriere).

60  
61 it is a cut-point more and more important in forensic sciences in  
62 order to decide if a children can be charged legally, the purpose  
63 of the present study was to examine the open apices of the teeth  
64 in discriminating between children who are or are not 14 years  
65 of age or older. If a child is younger than 12 or older than 16  
66 years of age there are many different and reliable techniques to  
67 assess that he/she is or is not younger than 14 years of age. For  
68 instance, taking into account the results in [10,2], it is easy to  
69 assess for a children older than 16 years that he/she is older than  
70 14 years of age. Furthermore, using the regression formula  
71 found in Cameriere et al. [11] it is possible to estimate that  
72 when all apices are closed a child is almost sure older than 12  
73 years (the probability that a child is younger than 12 years is  
74 less than 1%). Consequently, we considered children aged from  
12 to 16 years old as our target population.

76 **2. Materials and methods**

77 **2.1. Subjects and materials**

78 Orthopantomograms (OPGs) of 447 persons aged between 12 and 16 years,  
79 of Italian, Croatian and Slovenian nationality, were evaluated (Tables 1 and 2).  
80 Subdivision according to sex was similar, with 47% females and 53% males.  
81 Selection criteria for inclusion of OPGs in this study were: Caucasian origin; all  
82 teeth on the right lower jaw present; no obvious dental pathology on panoramic  
83 radiology related to the right lower jaw, tilted third molar. Only 21 (5.9%) of the  
84 447 OPGs examined, in which these criteria were not satisfied, were excluded.

85 To discriminate between individuals who are or are not aged 14 years or  
86 more, we analysed the apical ends of the roots of the seven left permanent  
87 mandibular teeth of each individual. Briefly, for each individual, we considered  
88 the following measures: (1) number of the seven left permanent mandibular  
89 teeth with root development complete, apical ends of roots completely closed  
90 ( $N_0$ ) and dichotomous variable  $C$  (with  $C = 1$ , if all seven left permanent teeth  
91 had completely closed apices;  $C = 0$ , if at least one tooth had its apices not  
92 completely closed) and (2) third molar maturity index ( $D_{3M}$ ), i.e., sum of the  
93 distances between the inner sides of the two open apices when roots were  
94 developed; otherwise,  $D_{3M}$  was obtained dividing crown length by tooth length  
95 of highest cusp [11].

96 Dental maturity was evaluated using the third molar maturity index  $D_{3M}$  and  
97 the dichotomous variable  $C$ .

98 **2.2. Statistical analysis**

99 All measurements were carried out by two observers. In order to evaluate  
100 intra- and inter-observer reliability, the two observers made repeated measures  
101 of 30 OPGs at an interval of 2 weeks.

102 The intra- and inter-observer reproducibility of the sum of the distances  
103 between the inner sides of the two open apices divided by the tooth length ( $D_{3M}$ )  
104 was studied using the concordance correlation coefficient,  $\rho_c$ , and  $\kappa$  statistics  
105 were used to measure the intra- and inter-observer reproducibility of the number  
106 of the seven right permanent mandibular teeth with root development complete  
107 ( $N_0$ ).

108 Using individual age as a dichotomous response variable ( $F = 1$  if an  
109 individual is at least 14 years of age,  $F = 0$  otherwise), and gender, nationality,

Table 1  
Countries distribution of the sample

Country	Females	Males	Total
Italy	84	85	169
Croatia	73	73	146
Slovenia	54	78	132
Total	211	236	447

Table 2  
Age and sex distribution of the sample

Years	Females	Males	Total
12	43	66	109
13	46	41	87
14	53	50	103
15	30	33	63
16	40	45	85
Total	212	235	447

110  $C$ , and  $D_{3M}$  as predictor variables, we derived a generalized linear model to  
111 predict whether an individual is older ( $F = 1$ ) or younger ( $F = 0$ ) than 14 years  
112 of age by using a logistic model as link function.

113 The predictive accuracy of the model was assessed by the determination of  
114 receiver operating characteristic curve (ROC curve).

115 All the significant variables were used to test the medico-legal question of  
116 whether an individual is older or younger than 14 years of age. The test was  
117 performed identifying a threshold (cut-off) that can be used to assign an  
118 individual to the population of the younger ( $T = 0$ ) or older ( $T = 1$ ) than 14  
119 years of age.

120 Sensitivity  $p_1$  of test (i.e., the proportion of children equal to or older than 14  
121 years of age, which verifies event  $T = 1$ ) was evaluated, and also its specificity,  
122  $p_2$  (i.e., the proportion of individuals younger than 14 years of age that verify the  
123 event  $T = 0$ ).

124 Open apices in teeth may help in discriminating between children who are or  
125 are not aged 14 years or more, by using the post-test probability of being 14  
126 years of age or more (i.e., the proportion of individuals aged 14 or over in whom  
127 event  $F = 1$  is verified). According to Bayes' theorem, post-test probability may  
be written as:

$$p = \frac{p_1 p_0}{p_1 p_0 + (1 - p_2)(1 - p_0)} \tag{2.1}$$

128 where  $p$  is post-test probability and  $p_0$  is the probability that a child is equal to or  
129 older than 14 years, given that he/she is aged between 12 and 16 years, which  
130 represents our target population. This probability,  $p_0$ , was evaluated using the  
131 data obtained from the statistical offices of Slovenia, Croatia and Italy [12-14].  
132 Since sensitivity and specificity, the determinants of post-test probability of  
133 being aged 14 years or more, were unknown probabilities, they were estimated  
134 using our sample subjects. Consequently, post-test probability  $p$  in Eq. (2.1)  
135 became a sample statistic subjected to random error. Thus, confidence intervals  
136 were used to describe its uncertainty.

137 The expression of the asymptotic  $(1 - \alpha)$  per cent confidence interval for the  
138 post-test probability estimate may be written in terms of the estimates of adult  
139 subjects, sensitivity, specificity and their corresponding sample size, as follows:

$$\left[ \frac{1}{1 + p \exp(z_{\alpha/2} \sqrt{V})}, \frac{1}{1 + p \exp(-z_{\alpha/2} \sqrt{V})} \right], \tag{2.2}$$

140 with

$$V = \frac{1 - p_1}{n_1 p_1} + \frac{p_2}{n_2 (1 - p_2)} \tag{2.3}$$

141 where  $n_0$  is the sample size, and  $n_1$  and  $n_2$  are the numbers of individuals who are  
142 or are not aged 14 years or more. Statistical analysis of data and related graphs  
143 was carried out with S-PLUS 6 statistical program (S-PLUS® 6.1 for Windows  
144 Professional Edition Release 1) and the Microsoft Excel® program. The  
145 significance level was set at 5%.

146 **3. Results**

147 For the number of the seven right permanent mandibular  
148 teeth with root development complete ( $N_0$ ), we did not observe  
149 any disagreement between two measurements made by the  
150 same observer, i.e.,  $\kappa = 1$ .

Inter-observer reproducibility of  $N_0$  was good with Cohen's  $\kappa$  statistics ( $\pm$ S.D.) at  $\kappa = 0.93 \pm 0.07$ , indicating no significant inter-observer differences.

As regards the reproducibility of  $D_{3M}$  measurements (sum of distances between inner sides of two open apices divided by tooth length), the estimated concordance correlation coefficient ( $\pm$ S.D.) was  $\rho_c = 0.966 \pm 0.0005$  for observer 1,  $\rho_c = 0.964 \pm 0.0035$  for observer 2, and  $\rho_c = 0.956 \pm 0.0076$ , when the measures of both observers were compared.

Inter-observer reproducibility of  $D_{3M}$  did not reveal significant intra- or inter-observer effects, indicating substantial homogeneity of evaluation between operators.

From the data at our disposal, it is inferred that, in 5.9% of the subjects examined, the third molar on the right lower jaw was not present.

For the remaining 94.1% of the data, we studied the extent to which the age of 14 years or more of an individual ( $F = 1$ ) is related to the maturation degree of the third molar ( $D_{3M}$ ), the dichotomous variable  $C$ , gender (1 for male and 0 for female) and nationality of the children.

Let  $p = P(F = 1)$  the probability that the an individual is at least 14 years of age, we modeled the dependence of this probability on  $D_{3M}$ ,  $C$ , gender and nationality using a linear logistic model:

$$\text{logit}(p) = b_0 + b_1 \text{nationality} + b_2 \text{gender} + b_3 C + b_4 D_{3M}. \quad (3.1)$$

To examine the effect of including one of the four factors in, or excluding it from the model, we considered the difference in deviance between two nested models (Table 3).

The change in deviance on adding the variables nationality and gender to a model that includes a constant term alone (null model) was not significant.

Instead, when  $C$  or  $D_{3M}$  were added to the null model, the deviance was reduced by highly significant amounts ( $p < 0.001$ ).

In summary, the probability that an individual is aged 14 years or more depends both on the dichotomous variable  $C$  which is related to the number of the seven left permanent mandibular teeth with root development complete and to the maturation degree of the third molar  $D_{3M}$ , but it does not significantly depend on gender and nationality. Hence Eq. (3.1) can be rewritten as:

$$p = \frac{1}{1 + e^{-(b_0 + b_1 C + b_2 D_{3M})}}. \quad (3.2)$$

Table 3  
Deviance on fitting the considered linear models to the data

	d.f.	Dev. resid.	d.f.	Deviance	$p$
Null	-	-	425	582.6	-
Nationality	1	1.6	424	581.0	0.20
Gender	1	0.1	423	580.9	0.71
$C$	1	323.0	422	257.8	<0.001
$D_{3M}$	1	16.3	421	241.5	<0.001

Terms were added sequentially (first to last).

Table 4  
Parameter estimates for logistic model (3.2)

Parameter	Value	S.E.	$t$ -Value
Intercept	0.308	0.530	0.581
$C$	4.233	0.367	11.527
$D_{3M}$	-2.190	0.544	-4.03

The maximum likelihood estimates of the model parameters (Table 4) evaluated the probability that an individual was equal to or older than 14 years of age,  $p$ , given the values of the factor  $C$  and covariate  $D_{3M}$  through the logistic model (3.2).

The predictive accuracy of Eq. (3.2) and its discrimination capacity was also assessed by determining the ROC curve by classification matrices for different levels of predicted probability that an individual is of age. The resulting ROC curve (Fig. 1) has an area under the curve ( $\pm$ S.D.) of  $0.814 \pm 0.021$ .

To test the legal question of whether an individual is older or younger than 14 years of age, a procedure had to be identified, such that an individual is assigned to the population of the younger than 14 years of age if the test is resulted negative ( $T = 0$ ) and to the older population if the test is resulted positive ( $T = 1$ ).

For forensic purposes, it is important that the test shows a low proportion of individuals younger than 14 years of age whose test is resulted positive ( $T = 1$ ), and so it seemed appropriate to pay more attention to the chance of a false positive than to that of a false negative.

On these grounds, we established that an individual is considered equal or older than 14 years of age (the test is positive,  $T = 1$ ) if  $C = 1$  and  $D_{3M}$  is lower than  $D_{3M}^* = 1.1$ ; otherwise an individual is considered younger than 14 years of age (the test is negative,  $T = 0$ ).

The sensitivity of this test (the proportion of individuals being older or equal to 14 years of age whose test is positive) was 81%, and its specificity (the proportion of individuals younger than 14 years of age whose test is negative) was 95%.




Fig. 1. Receiver operating characteristic curve for "14 years of age or older" status.

Table 5  
Classification table describing discrimination performance of the test

	Age		Total
	<14	≥14	
T = 0	175	46	221
T = 1	9	196	205
Total	184	242	426

Table 6  
Distribution of the age for gender and test

Gender	T	Age						Total
		12	13	14	15	16	17	
F	0	0.21	0.18	0.04	0.03	0.03	0.00	0.47
	1	0.00	0.03	0.22	0.11	0.12	0.05	0.53
Total F		0.21	0.21	0.26	0.13	0.15	0.05	1.00
M	0	0.28	0.14	0.06	0.02	0.01	0.00	0.52
	1	0.00	0.02	0.15	0.13	0.14	0.04	0.48
Total M		0.28	0.16	0.21	0.15	0.15	0.04	1.00

The proportion of individuals with correct classifications was 87% (Table 5).

In the sample, estimated post-test probability  $p$  was 0.96, with a 95% confidence interval, CI = (0.93, 0.98). Hence, the probability that a subject positive on the test ( $T = 1$ ) was equal or older than 14 years of age was 96%. Consequently, the test yielded only 4% of false positives.

When subjects of 12 years of age were examined, using this test, no subjects were estimated as older than or equal to 14 years of age. In addition, when subjects of 13 years of age were examined, the test estimated only 2% of both males and females as older than 14 (Table 6). Furthermore, when subjects of 16 years of age were examined, none of them were estimated younger than 14 years of age.

#### 4. Discussion

The need for effective and reliable scientific methods to determine age, particularly adult and over age of 14 years old, within a specific population has become increasingly important in resolving court cases. Since the methods usually applied for dental age estimation guarantee an error in estimated age of less than 2 years [8,15,16], to estimate the post-test probability and prevalence of subjects older than 14, we chose young people aged between 12 and 16 years old as a target population.

Our results showed that the test is not significantly dependent on the nationality (Croatian, Italian and Slovenian) of the children neither to their gender while it depends on the maturation degree of the teeth.

In this paper, our test estimates that a subject is older than 14 years of age if all the teeth, except the third molar, have closed apices (are fully grown) and the maturation degree of the third molar,  $D_{3M}$ , is equal or lesser than 1.1.

When the suggested test was applied, the percentage of false negatives was 19% and the percentage of false positives was

5%. From a forensic point of view, the small percentage of false positives is particularly important, because it is a more serious error to consider a subject younger than 14 as chargeable than the error which does not consider a subject older than 14 as chargeable.

Our results confirmed that, if the root apices of the seven teeth in the right lower jaw of a child are completely closed, and the ratio of the sum of third molar root apices divided by tooth length is lower than 1.1, then there is a high probability that the subject is indeed at least 14 years of age. In fact, the estimated probability that a child with  $C = 1$  and  $D_{3M} \leq 1.1$  has reached 14 years of age is  $p = 0.96$ .

In Cameriere et al. [17] we analysed a technique to assess biological growth and age in children and adolescents using the wrist/hand area method. Ossification of the carpals showed good agreement with chronological age, and their mineralization lasts until the age of approximately 14. For this reason, analysis is in progress to assess the age of boys and girls in the 12–16 age bracket using a combination of carpal bone and tooth growth information.

#### Acknowledgement

This work was supported by a grant from the Polytechnic University of Marche.

#### References

- [1] P.M. Garamendi, M.I. Landa, J. Ballesteros, M.A. Solano, Reliability of the methods applied to assess age minority in living subjects around 18 years old. A survey on a Moroccan origin population, *Forensic Sci. Int.* 154 (2005) 3–12.
- [2] A. De Salvia, C. Calzetta, M. Orrico, D. De Leo, Third mandibular molar radiological development as an indicator of chronological age in a European population, *Forensic Sci. Int.* 146 (2004) S9–S12.
- [3] N. Chaillat, A. Demirjian, Dental maturity in South France: a comparison between Demirjian's method and polynomial functions, *J. Forensic Sci.* 49 (2004) 1059–1066.
- [4] A. Olze, A. Mahlow, S. Schmidt, K.D. Wernecke, G. Geserick, A. Schemling, Combined determination of selected radiological and morphological variables relevant for dental age estimation of young adults, *Homo* 56 (2005) 133–140.
- [5] H.M. Liversidge, F. Lyons, M.P. Hector, The accuracy of three methods of age estimation using radiographic measurements of developing teeth, *Forensic Sci. Int.* 131 (2003) 22–29.
- [6] J.M. Tanner, M.J.R. Healy, H. Goldstein, N. Cameron, Assessment of Skeletal Maturity and Prediction of Adult Height (TW3 Method), W.D. Saunders, London, 2001.
- [7] A.F. Roche, W. Cameron Chumlea, D. Thissen, Assessing the Skeletal Maturity of the Hand–Wrist: Fels Method, Charles C. Thomas, Springfield, 1988.
- [8] A. Demirjian, H. Goldstein, J.M. Tanner, A new system of dental age assessment, *Hum. Biol.* 45 (1973) 221–227.
- [9] G. Willems, A review of the most commonly used dental age estimation techniques, *J. Forensic Odontostomatol.* 19 (2001) 9–17.
- [10] H.H. Mincer, E.F. Harris, H.E. Berryman, The A.B.F.O. study of third molar development and its use as an estimator of chronological age, *J. Forensic Sci.* 3 (1993) 379–390.
- [11] R. Cameriere, L. Ferrante, M. Cingolani, Age estimation in children by measurement of open apices in teeth, *Int. J. Legal Med.* 120 (2006) 49–52.
- [12] Statistical Office of the Republic of Slovenia, [http://www.stat.si/letopis/index\\_letopis\\_en.asptable4.4](http://www.stat.si/letopis/index_letopis_en.asptable4.4).

319  
320  
321  
322  
323  
324  
325

- [13] Republic of Croatia central Bureau of Statistics, <http://www.dzs.hr/ljetopis/LjFrameE.htm>.
- [14] Istituto Nazionale di Statistica (ISTAT), <http://demo.istat.it/pop2004/index.html>.
- [15] C.S. Farah, D.R. Booth, S.C. Knott, Dental maturity of children in Perth, Western Australia, and its application in forensic age estimation, *J. Clin. Forensic Med.* 6 (1999) 14-18.

- [16] A. Teivens, H. Mornstad, A comparison between dental maturity rate in the Swedish and Korean populations using a modified Demirjian method, *J. Forensic Odontostomatol.* 19 (2001) 31-35.
- [17] R. Cameriere, L. Ferrante, D. Mirtella, M. Cingolani, Carpals and epiphyses of radius and ulna as age indicators, *Int. J. Legal Med.* 7 (2005) 1-4.

325  
326  
327  
328  
329  
330  
331  
331

UNCORRECTED PROOF